145 The West Mall, P.O. Box 110 U
Toronto, ON M8Z 5M4

PRIOR AUTHORIZATION PROGRAM FORM

Jakavi ® (Ruxolitinib)

Instructions:

1. Section 1 to be completed by Plan Member / Patient
2. Section 2 to be completed by Physician (expenses incurred by the completion of this form is at plan member's expense)
3. Section 3 to be completed by Pharmacist

Please fax completed form to NexgenRx Formulary Management to 1-877-639-4369 or mail to 145 The West Mall P.O. Box 110 U, Toronto, Ontario,M8Z-5M4

Section 1: To be completed by Patient

Member's Name: (Last , First)	Card ID Number:	
Patient's Name: (Last, First)	Patient's Date of Birth (dd/mm/yy)	Relationship to Member (please circle) Employee Spouse Dependent

PATIENT CODE

Results of this request to be communicated to:

☐ Pharmacy ☐ Patient / Legal Guardian Named Below ☐ email:

I authorize NexgenRx Inc. (a) to use the personal information disclosed on this form, and any other personal information known to NexgenRx Inc. regarding the above-named patient, for the purpose of assessing this prior authorization request and any related claim and administering the benefit plan under which any such claim is made, and (b) to contact, and to obtain any such personal information from and to disclose any such personal information to, any physician, pharmacist or other health care professional having knowledge of such patient's health relevant to this request and any related claim.

Patient / Legal Guardian Name: _____ Telephone Number: _____

Signature of Patient/ Legal Guardian: _____ Date (dd/mm/yy):_____

Section 2: To be completed by Physician

Drug Name & Strength:	DIN:	Dosage Instructions:

Eligibility for drug coverage is dependent upon the patient meeting at least one of the qualifying criteria listed below. If the patient has another drug plan, prior authorization may cover some or all of the excess not paid by the primary plan.

Please indicate if the patient meets the following qualifying criteria for drug coverage:

☐ Jakavi is indicated for the treatment of splenomegaly and/or its associated symptoms in adult patients with primary myelofibrosis (also known as chronic idiopathic myelofibrosis), post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis.

☐ Jakavi is indicated in the control of hematocrit in adult patients with polycythemia vera (PV) resistant to or intolerant of a cytoreductive agent.

Physician Name: (Last, First)	License Number:
Address: (Street, City, Province, Postal Code)	Telephone Number : ( ) - Fax Number: ( ) -
Signature of Physician:	Date (dd/mm/yy)

Section 3: To be completed by Pharmacist

Pharmacy Name:	Provider Number:
Pharmacy Address: (Street, City, Province, Postal Code)	Telephone Number : ( ) - Fax Number: ( ) -
Signature of Pharmacist: _____ Pharmacist's Name: (Print Last, First) _____	

Internal Office Use Only:

Date Received:			Date Approved & By: